

2  
paper number

REQUEST FOR ACCESS

Date: 7-3-96  
Serial Number: 07/310252  
Filing Date: 2-13-89  
Applicants: QUEEN ET AL.

Sir.

Request is hereby respectfully made for access to the file history of the following  
abandoned application referred to in U.S. patent number 5530101 or printed  
application number \_\_\_\_\_

Respectfully submitted.

WAYNE CROTEAU

-----  
OFFICIAL USE ONLY  
-----

PROCESSED BY

JUL 3 - 1996

FII

\_\_\_\_\_  
initials

FILE INFORMATION UNIT  
-----



US005530101A

**United States Patent** [19]  
**Queen et al.**

[11] **Patent Number:** **5,530,101**  
 [45] **Date of Patent:** **Jun. 25, 1996**

- [54] **HUMANIZED IMMUNOGLOBULINS**
- [75] **Inventors:** Cary L. Queen, Los Altos; Harold E. Selick, Belmont, both of Calif.
- [73] **Assignee:** Protein Design Labs, Inc., Mountain View, Calif.
- [21] **Appl. No.:** 634,278
- [22] **Filed:** Dec. 19, 1990

**Related U.S. Application Data**

- [63] Continuation-in-part of Ser. No. 590,274, Sep. 28, 1990, abandoned, and a continuation-in-part of Ser. No. 310,252, Feb. 13, 1989, abandoned, which is a continuation-in-part of Ser. No. 290,975, Dec. 28, 1988, abandoned.
- [51] **Int. Cl.<sup>6</sup>** ..... **A61K 39/395; C07K 16/28**
- [52] **U.S. Cl.** ..... **530/387.3; 530/387.1; 530/388.22; 424/133.1; 424/143.1**
- [58] **Field of Search** ..... **424/85.8, 133.1, 424/143.1; 530/387, 388.22, 387.1, 387.3**

**References Cited**

**U.S. PATENT DOCUMENTS**

4,816,397	3/1989	Boss et al.	435/68
4,816,567	3/1989	Cabilly et al.	530/387
4,867,973	9/1989	Geers et al.	
5,225,539	7/1993	Winter	

**FOREIGN PATENT DOCUMENTS**

0171496	2/1986	European Pat. Off.	
0173494	3/1986	European Pat. Off.	
0184187	6/1986	European Pat. Off.	
0239400	9/1987	European Pat. Off.	
0266663	6/1988	European Pat. Off.	
2188941	10/1987	United Kingdom	
WO86/05513	9/1986	WIPO	
WO87/02671	5/1987	WIPO	
WO89/01783	3/1989	WIPO	

**OTHER PUBLICATIONS**

Vitteta et al., "Redesigning Nature's Poisons to Create Anti-Tumor Reagents," *Science* 238:1098-1104 (1987).  
 Ellison et al., "The nucleotide sequence of a human immunoglobulin C(gamma)<sub>1</sub> gene", *Nucleic Acids Res.* 10:4071-(1982).  
 Hieter et al., "Cloned Human and Mouse Kappa Immunoglobulin Constant and J Region Genes Conserve homology in Functional Segments", *Cell* 22:197-207 (1980).  
 Sharon et al., "Expression of a V<sub>H</sub>C<sub>K</sub> chimaeric protein in mouse myeloma cells", *Nature* 309:364-367 (1984).  
 Takeda et al., "Construction of chimaeric processed immunoglobulin genes containing mouse variable and human constant region sequences", *Nature* 314:452-454 (1985).  
 Tan et al., "A Human-Mouse Chimeric Immunoglobulin Gene with a Human Variable Region is Expressed in Mouse Myeloma Cells", *J. Immunol.* 135:3564-3567 (1985).  
 Morrison et al., "Chimeric human antibody molecules: Mouse antigen-binding domains with human constant region domains," *Proc. Natl. Acad. Sci.* 81:6851-6859 (1984).  
 Boulianne et al., "Production of functional chimeric mouse/human antibody," *Nature* 312:643-646 (1984).

Neuberger et al., "A hapten-specific chimeric IgE antibody with human physiological effector function," *Nature* 314:268-270 (1985).  
 Morrison, S. L., "Transfectomas Provide Novel Chimeric Antibodies," *Science* 229:1202-1207 (1985).  
 Sahagan et al., "A Genetically Engineered Murine/Human Chimeric Antibody Retains Specificity for Human Tumor-Associated Antigen", *J. Immunol.* 137:1066-1074 (1986).  
 Liu et al., "Expression of mouse::human immunoglobulin heavy-chain cDNA in lymphoid cells", *Gene* 54:33-40 (1987).  
 Better et al., "Escherichia coli Secretion of an Active Chimeric Antibody Fragment", *Science* 240:1041-1043 (1988).  
 Waldmann, T. A., "The Structure, Function, and Expression of Interleukin-2 Receptors on Normal and Malignant Lymphocytes," *Science* 232:727-732 (1986).  
 Leonard et al., "The human receptor for T-cell growth factor," *J. Biol. Chem.* 260:1872-1880 (1985).  
 Farrar, J., "The biochemistry, biology, and role of interleukin-2 in the induction of cytotoxic T cell and antibody-forming B cell receptors," *Immunol. Rev.* 63:129-166 (1982).  
 Greene et al., "Growth of Human T Lymphocytes: An Analysis of Interleukin 2 and Its Cellular receptor", in *Progress in Hematology XIV*, E. Brown ed., Grune and Stratton, New York (1986) pp. 283-301.  
 Verhoyen et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity", *Science* 239:1534-1536 (1988).  
 Jones et al., "Replacing the complementarity-determining regions in a human antibody with those from a mouse", *Nature* 321:522-525 (1986).  
 Hale et al., "Remission Induction in Non-Hodgkin Lymphoma with Reshaped Human Monoclonal Antibody CAMPATH-1H", *Lancet* Dec. 17, 1988, pp. 1394-1399.  
 Chothia, C. and A. M. Lesk, "Canonical Structures for the Hypervariable Regions of Immunoglobulins", *J. Mol. Biol.* 196:901-917 (1987).

(List continued on next page.)

**Primary Examiner**—Lila Feisce  
**Attorney, Agent, or Firm**—Townsend and Townsend and Crew

[57] **ABSTRACT**  
 Novel methods for producing, and compositions of, humanized immunoglobulins having one or more complementarity determining regions (CDR's) and possible additional amino acids from a donor immunoglobulin and a framework region from an accepting human immunoglobulin are provided. Each humanized immunoglobulin chain will usually comprise, in addition to the CDR's, amino acids from the donor immunoglobulin framework that are, e.g., capable of interacting with the CDR's to effect binding affinity, such as one or more amino acids which are immediately adjacent to a CDR in the donor immunoglobulin or those within about 3 Å as predicted by molecular modeling. The heavy and light chains may each be designed by using any one or all of various position criteria. When combined into an intact antibody, the humanized immunoglobulins of the present invention will be substantially non-immunogenic in humans and retain substantially the same affinity as the donor immunoglobulin to the antigen, such as a protein or other compound containing an epitope.